Criteria-Based Consultation Prescribing Program CRITERIA FOR DRUG COVERAGE

Insulin glargine (Basaglar Kwikpen, Basaglar Tempo Pen)

Notes:

- Quantity Limits: No
- * Intolerance excludes adverse drug reactions that are expected, mild in nature, resolve with continued treatment, and do not require medication discontinuation
- ^ Insulinopenia is defined fasting c-peptide less than or equal to 0.88 ng/mL (or 1.6 ng/mL in patients with creatinine clearance less than 50 mL/min) with a concurrent blood glucose of 70-225 mg/dL

<u>Initiation (new start) criteria</u>: Non-formulary **insulin glargine (Basaglar Kwikpen, Basaglar Tempo Pen)** will be covered on the prescription drug benefit when the following criteria are met:

 The patient has a documented allergic reaction to an inactive ingredient in Lantus, or its unbranded biologic, and Semglee, or its unbranded biologic, not present in Basaglar

-AND-

Meets one of the following criteria:

- Use in patients with type 1 diabetes mellitus as basal insulin
- Use in patients with any type of diabetes age 19 or younger
- Use in patients with type 2 diabetes mellitus AND insulinopenia[^]
- Use in patients with type 2 diabetes mellitus AND documented allergy or intolerance* to NPH insulin
- Use in patients with type 2 diabetes mellitus that experience recurrent nocturnal hypoglycemia (low blood sugar at night) with bedtime NPH insulin dosing defined as: 3 or more episodes of nocturnal CBG (capillary blood glucose) less than 70 over the preceding 30 days that persists despite NPH insulin dose reduction
- For patients on 70/30 insulin, trial of NPH insulin (dosed am and bedtime) and Regular insulin (dosed breakfast and dinner) where the bedtime dose of NPH insulin resulted in recurrent hypoglycemia as defined above
- Use in patients with type 2 diabetes mellitus on NPH insulin that experience any
 episode of severe hypoglycemia defined as: hypoglycemia resulting in seizures, loss
 of consciousness, episode necessitating assistance from someone else, EMT
 (emergency medical technician), and/or use of glucagon (medication used to raise
 the concentration of glucose in the blood)

kp.org

Revised: 7/14/22 Effective: 9/1/22 All plans offered and underwritten by Kaiser Foundation Health Plan of the Northwest




Criteria-Based Consultation Prescribing Program CRITERIA FOR DRUG COVERAGE

Insulin glargine (Basaglar Kwikpen, Basaglar Tempo Pen)

 Use in patient with type 2 diabetes mellitus that require long-acting insulin due to work (night shift work where hours of sleep are significantly and repeatedly varied over time, frequent time-zone traveler)

kp.org

Revised: 7/14/22 Effective: 9/1/22





All plans offered and underwritten by