

Criteria-Based Consultation Prescribing Program

CRITERIA FOR DRUG COVERAGE

Ticagrelor (Brilinta)

Notes:

- Quantity Limits: No
- * Examples of clopidogrel failure includes acute coronary syndrome, stent thrombosis, or stroke while on clopidogrel

Initiation (new start) criteria: Formulary **ticagrelor (Brilinta)** will be covered on the prescription drug benefit when the following criteria are met:

For patients with documented allergy, intolerance, or treatment failure with clopidogrel, formulary **ticagrelor (Brilinta)** will be covered on the prescription drug benefit when the following criteria are met:

- Prescribed by or in consultation with cardiology, neurology, or vascular surgery
- *AND* -
- Allergy or intolerance to clopidogrel therapy
- *OR* -
- Failed clopidogrel therapy*

For patients WITHOUT documented allergy, intolerance, or treatment failure with clopidogrel, formulary **ticagrelor (Brilinta)** will be covered on the prescription drug benefit when the following criteria are met:

- Prescribed by or in consultation with interventional cardiologist
- *AND* -
- Used in patients with stent placement in the setting of acute myocardial infarction
- *AND* -
- Receiving concomitant aspirin 81 mg daily (not to exceed 100 mg daily)
- *AND* -
- Does not have any of the following exclusions:
 - History of intracranial hemorrhage
 - Concomitant use of strong CYP3A4 inducers or inhibitors
 - Concomitant oral anticoagulant therapy
 - Severe hepatic impairment

Criteria for current Kaiser Permanente members already taking the medication who have not been reviewed previously: Formulary **ticagrelor (Brilinta)** will be covered on the prescription drug benefit when the following criteria are met:

kp.org

Revised: 08/10/23
Effective: 10/05/23

All plans offered and underwritten by
Kaiser Foundation Health Plan of the Northwest

Criteria-Based Consultation Prescribing Program

CRITERIA FOR DRUG COVERAGE

Ticagrelor (Brilinta)

For patients with documented allergy, intolerance, or treatment failure with clopidogrel, formulary **ticagrelor (Brilinta)** will be covered on the prescription drug benefit when the following criteria are met:

- Prescribed by or in consultation with cardiology, neurology, or vascular surgery
- *AND* -
- Allergy or intolerance to clopidogrel therapy
- *OR* -
- Failed clopidogrel therapy*

For patients WITHOUT documented allergy, intolerance, or treatment failure with clopidogrel, formulary **ticagrelor (Brilinta)** will be covered on the prescription drug benefit when the following criteria are met:

- Prescribed by or in consultation with interventional cardiologist
- *AND* -
- Used in patients with stent placement in the setting of acute myocardial infarction
- *AND* -
- Receiving concomitant aspirin 81 mg daily (not to exceed 100 mg daily)
- *AND* -
- Does not have any of the following exclusions:
 - History of intracranial hemorrhage
 - Concomitant use of strong CYP3A4 inducers or inhibitors
 - Concomitant oral anticoagulant therapy
 - Severe hepatic impairment

Criteria for new members entering Kaiser Permanente already taking the medication who have not been reviewed previously: Formulary **ticagrelor (Brilinta)** will be covered on the prescription drug benefit when the following criteria are met:

For patients with documented allergy, intolerance, or treatment failure with clopidogrel, formulary **ticagrelor (Brilinta)** will be covered on the prescription drug benefit when the following criteria are met:

- Prescribed by or in consultation with cardiology, neurology, or vascular surgery
- *AND* -
- Allergy or intolerance to clopidogrel therapy
- *OR* -
- Failed clopidogrel therapy*

kp.org

Revised: 08/10/23
Effective: 10/05/23

All plans offered and underwritten by
Kaiser Foundation Health Plan of the Northwest

Criteria-Based Consultation Prescribing Program

CRITERIA FOR DRUG COVERAGE

Ticagrelor (Brilinta)

For patients WITHOUT documented allergy, intolerance, or treatment failure with clopidogrel, formulary **ticagrelor (Brilinta)** will be covered on the prescription drug benefit when the following criteria are met:

- Prescribed by or in consultation with interventional cardiologist
- *AND* -
- Used in patients with stent placement in the setting of acute myocardial infarction
- *AND* -
- Receiving concomitant aspirin 81 mg daily (not to exceed 100 mg daily)
- *AND* -
- Does not have any of the following exclusions:
 - History of intracranial hemorrhage
 - Concomitant use of strong CYP3A4 inducers or inhibitors
 - Concomitant oral anticoagulant therapy
 - Severe hepatic impairment