## Criteria-Based Consultation Prescribing Program CRITERIA FOR DRUG COVERAGE

## Insulin detemir pen (Levemir FlexPen)

#### Notes:

\* Intolerance excludes adverse drug reactions that are expected, mild in nature, resolve with continued treatment, and do not require medication discontinuation

^ Insulinopenia is defined fasting c-peptide less than or equal to 0.88 ng/mL (or 1.6 ng/mL in patients with creatinine clearance less than 50 mL/min) with a concurrent blood glucose of 70-225 mg/dL

Non-formulary **insulin detemir pen (Levemir FlexPen)** will be covered on the prescription drug benefit when the following criteria are met:

Documented allergy or intolerance\* to insulin glargine AND insulin degludec

#### -AND-

#### Meets one of the following criteria:

- Use in patients with type 1 diabetes mellitus as basal insulin
- Use in patients with any type of diabetes age 19 or younger
- Use in patients with type 2 diabetes mellitus AND insulinopenia<sup>^</sup>
- Use in patients with type 2 diabetes mellitus AND documented allergy or intolerance\* to NPH insulin
- Use in patients with type 2 diabetes mellitus that experience recurrent nocturnal hypoglycemia (low blood sugar at night) with bedtime NPH insulin dosing defined as: 3 or more episodes of nocturnal CBG (capillary blood glucose at night) less than 70 over the preceding 30 days that persists despite NPH insulin dose reduction
  - For patients on 70/30 insulin, trial of NPH insulin (dosed am and bedtime) and Regular insulin (dosed breakfast and dinner) where the bedtime dose of NPH insulin resulted in recurrent hypoglycemia as defined above
- Use in patients with type 2 diabetes mellitus on NPH insulin that experience any
  episode of severe hypoglycemia defined as: hypoglycemia resulting in seizures,
  loss of consciousness, episode necessitating assistance from someone else, EMT
  (emergency medical technician), and/or use of glucagon (medication used to raise
  the concentration of glucose in the blood)
- Use in patient with type 2 diabetes mellitus that require long-acting insulin due to work (night shift work where hours of sleep are significantly and repeatedly varied over time, frequent time-zone traveler)

#### -AND-

kp.org

Revised: 06/08/23 Effective: 08/03/23 All plans offered and underwritten by Kaiser Foundation Health Plan of the Northwest



# Criteria-Based Consultation Prescribing Program CRITERIA FOR DRUG COVERAGE


### Insulin detemir pen (Levemir FlexPen)

### Meets one of the following criteria:

- Unable to draw up insulin accurately from a vial with a syringe due to young age, visual impairment, physical disabilities (i.e., amputations, tremors/Parkinson's disease, rheumatoid arthritis), or history of IV drug use
- Requires small doses of insulin (less than 5 units/dose) or stabilized on smaller insulin dose (generally less than 30 units per day or less than 1000 units per month)
- Pediatric patient who is required to use such devices by school

kp.org

Revised: 06/08/23 Effective: 08/03/23





All plans offered and underwritten by