

Criteria-Based Consultation Prescribing Program

CRITERIA FOR DRUG COVERAGE

Ertugliflozin/ Metformin (Segluromet)

Initiation (new start) criteria: Non-formulary **ertugliflozin/metformin (Segluromet)** will be covered on the prescription drug benefit when the following criteria are met:

- Diagnosis of type 2 diabetes
- Intolerance* to SGLT-2 inhibitors empagliflozin (Jardiance), canagliflozin (Invokana) AND dapagliflozin (Farxiga)
- Recent HbA1c less than 10%
- On maximally tolerated metformin dose (dose appropriate per renal function) or allergy or intolerance* to metformin (includes both metformin IR and XR)
- Diagnosis of one or more of the following conditions:
 1. Atherosclerotic Cardiovascular Disease (ASCVD)**
 2. Urine albumin/creatinine ratio greater than 300 mg/g **AND** GFR greater than 59 mL/min (Do not initiate for GFR less than 60 mL/min)
 - And on maximally tolerated dose or patient has an allergy or intolerance* to ACE/ARB
 3. Heart Failure

* Intolerance excludes adverse drug reactions that are expected, mild in nature, resolve with continued treatment, and do not require medication discontinuation

** ASCVD refers to the following conditions:

- Coronary heart disease (CHD, CAD), such as myocardial infarction, angina, and coronary artery stenosis > 50%.
- Cerebrovascular disease, such as transient ischemic attack, ischemic stroke, and carotid artery stenosis > 50%.
- Peripheral artery disease, such as claudication.
- Aortic atherosclerotic disease, such as abdominal aortic aneurysm and descending thoracic aneurysm. (Patients with only incidental aortic atherosclerosis are not included)

Criteria-Based Consultation Prescribing Program

CRITERIA FOR DRUG COVERAGE

Ertugliflozin/ Metformin (Segluromet)

Criteria for members already taking the medication who have not been reviewed previously (e.g., new members): Non-formulary **ertugliflozin/metformin (Segluromet)** will be covered on the prescription drug benefit for when the following criteria are met:

- Diagnosis of type 2 diabetes
- Intolerance* to SGLT-2 inhibitors empagliflozin (Jardiance), canagliflozin (Invokana) AND dapagliflozin (Farxiga)
- On maximally tolerated metformin dose (dose appropriate per renal function) or allergy or intolerance* to metformin (includes both metformin IR and XR)
- Diagnosis of one or more of the following conditions:
 4. Atherosclerotic Cardiovascular Disease (ASCVD)**
 5. Urine albumin/creatinine ratio greater than 300 mg/g **AND** GFR greater than 59 mL/min (Do not initiate for GFR less than 60 mL/min)
 - And on maximally tolerated dose or patient has an allergy or intolerance* to ACE/ARB
 6. Heart Failure

* Intolerance excludes adverse drug reactions that are expected, mild in nature, resolve with continued treatment, and do not require medication discontinuation

** ASCVD refers to the following conditions:

- Coronary heart disease (CHD, CAD), such as myocardial infarction, angina, and coronary artery stenosis > 50%.
- Cerebrovascular disease, such as transient ischemic attack, ischemic stroke, and carotid artery stenosis > 50%.
- Peripheral artery disease, such as claudication.
- Aortic atherosclerotic disease, such as abdominal aortic aneurysm and descending thoracic aneurysm. (Patients with only incidental aortic atherosclerosis are not included)