



## Kaiser Foundation Health Plan - California

### Utilization Management (UM) Criteria for the Provision of Occupational and Physical Therapy Services-Commercial

#### Utilization Management Criteria Statement

This document includes criteria that support utilization management ("UM") of certain provider requested health care services. (Refer to the "UM Criteria References" section, below.)

UM occurs when a qualified practitioner other than the treating clinician reviews the treating clinician's request against UM criteria. The qualified practitioner is in the position to approve, deny, delay or modify the service request based on a determination of medical necessity. These criteria are consistent with professional standards of practice and are provided for your reference.

If you are in a treatment relationship with a member, then your clinical recommendations are not subject to these criteria. Your treatment recommendations are guided by your professional judgment and informed, where applicable, by clinical practice guidelines and clinical support tools found in the UM Criteria References section, below.

#### Principles

The Permanente Medical Group, Inc. ("TPMG") and Southern California Permanente Medical Group ("SCPMG") provides occupational and physical therapy services when medically necessary.

**Occupational therapy** provides task-oriented therapeutic activities and exercises designed to significantly improve, develop or restore physical functions lost or impaired; or to help an individual relearn basic daily living skills or compensatory techniques to improve the level of independence in the activities of daily living.

**Physical therapy** is a dynamic profession with an established theoretical and scientific base and widespread clinical applications in the restoration and promotion of optimal physical function. Physical therapists diagnose and manage movement dysfunction and enhance physical and functional abilities.

## Eligibility

KFHP is responsible for coordinating the provision of services with other entities, including but not limited to Regional Centers and County Mental Health plans, to ensure that KFHP and other entities are not providing duplicative services.

## Diagnostic and Treatment Indications for Medically Necessary Occupational and Physical Therapy Services

The following identifies the diagnostic and treatment indications for which occupational or physical therapy services may be medically necessary plus other considerations in determining medical necessity.

**Musculoskeletal Pathology or Dysfunction**, including limitations in joint range of motion and/or mobility, deterioration from previous function of muscle strength and/or decreased endurance, soft tissue dysfunction, alterations in postural control and alignment.

**Neuromuscular Pathology or Dysfunction**, including deterioration from previous function or significant delay of gross and/or fine motor coordination, alterations in tone- increased or decreased, deterioration from previous function or significant delay of motor planning skills, deterioration from previous function or significant delay of balance, loss of selective motor control, decrease in bilateral integration.

**Neurocognitive Pathology or Dysfunction**, including evaluation and treatment for sensory deficits when they impact overall health or cause significant impairment of function when there is a reasonable expectation that treatment will lead to improvement in health or function. Therapy is not provided for sensory disorders in the absence of a functional impairment.

**Pathology or Dysfunction of the Vascular System**, including primary or secondary lymphedema, edema and venous stasis.

**Pathology or Injury to Skin**, including burns and/or scars following injury or surgery, open wounds.

**Design of Maintenance Activities**, including physical exercise, drills, techniques that a patient performs outside of therapy or after any therapy has concluded.

**Assessments of Impairment**, including appropriate assessments as part of a multidisciplinary or interdisciplinary team of motor skills and/or activities of daily living impairment; appropriate assessments of post therapy functions and periodic reviews of appropriate maintenance activities.

**Significant delay**, when considering services for individuals with developmental delays and disorders shall take into account the following considerations:

- a. Whether the individual scores below the 7th percentile for the lower of his or her chronological age or developmental level (also calculated as 1.5 standard deviations below the member's expected mean) on a standardized test used in the evaluation of activities of daily living or motor skills; OR
- b. If the individual at any age is not able to participate in standardized testing (whether because of age or inability to understand or cooperate in the testing process), an occupational therapist or physical therapist designated by PMG has determined that the individual has a significant delay in activities of daily living or motor skills commensurate with consideration (a).

Occupational and physical therapy services are those that require the skills of licensed providers of physical therapy and occupational therapy, within such provider's scope of practice, and in accordance with law.

Occupational and physical therapy services are provided on an episodic basis.

Inpatient occupational and physical therapy services may be provided in the hospital, as appropriate.

Outpatient physical therapy and occupational therapy services are provided episodically in the physical therapy or occupational therapy medical office.

Home health occupational and physical therapy may be prescribed as part of a home health care plan and provided episodically in the home.

Note:

- Therapies, interventions and techniques for some behavioral and psychological symptoms of behavioral health care conditions, including developmental conditions, may be available from PMG behavioral health care providers or Occupational /Physical therapist

## **General principles governing intervention for occupational and physical therapies**

Occupational or physical therapy services can be indicated when, based upon evidence-based medicine, as determined by a physical or occupational therapist, as applicable, and a PMG physician who is competent to evaluate the specific clinical issues involved in the healthcare services requested, such that the patient will achieve:

- a. significant, measurable improvement in the patient's mobility or basic activities of daily living functions beyond what would have been expected through the normal course of maturation or healing without the services; OR
- b. significant, measurable reversal of deterioration from previous levels or significant measurable improvement beyond what would have been expected through the normal course of maturation or healing without the services. OR
- c. the skills of a qualified provider of OT or PT services are medically necessary to maintain function or prevent worsening of a condition.

Determinations regarding a "reasonable functional progress/maintenance/prevention of regression " referenced above shall be based upon evidence-based medicine and requires the skills of a qualified provider of OT/PT therapy services is medically necessary to preserve function or prevent worsening of a condition. Additionally, determinations regarding medical necessity, take into consideration whether, in the clinical judgment of a licensed occupational or physical therapist, as appropriate, an individual is an appropriate candidate for occupational or physical therapy based on the following: (a) ability to participate in and/or benefit from the therapy process; (b) adequate attention span, cooperation, and endurance to participate in active treatment in the therapy process; (c) demonstration of behavior conducive to engaging in the therapy process; and (d) reasonable ability to generalize the therapy from the episodic sessions.

### **Circumstances which Occupational or Physical Therapy Services May Not Be Medically Necessary**

Some of these circumstances in which physical or occupational therapy may not be medically necessary are described below:

- **Individualized Education Program (IEP) Development**--Services requested by educational systems for development of an IEP. However, reports describing medically necessary physical or occupational therapy services rendered according to these guidelines may also be provided to educational systems for the development of an IEP at the request of the patient.
- **Recreational therapy**-Activities targeting recreation, enhancement, or wellness; not addressing functional skills.
- **Activities that provide diversion or general motivation** (e.g. encouraging participation in sports, adaptive PE, weight loss)

- **Specialized job testing** for the sole purpose of determining a person's ability to perform a particular job
- **Services solely for palliative purposes**, including massage and whirlpool
- **Non-evidence-based techniques** such as Alexander technique, cranio-sacral therapy, rolfing.
- **Exercise Programs for healthy individuals**, including development and delivery of exercise programs; assisted walking
- **Maintenance programs.** Drills, techniques and exercises after completion of medically necessary physical or occupational therapy services to preserve the patient's present level of function and prevent regression of that function. Maintenance begins when the therapeutic goals of the treatment plan have been achieved or when no further functional progress is apparent or expected to occur. Maintenance does not require the skills of a qualified provider of physical or occupational therapy services. The patient is responsible for practicing learned drills, techniques and exercises to preserve his or her present level of function and prevent regression of that function. Maintenance includes ongoing supervision of independent exercise programs, supervision/observation of activities of daily living, and supervision of independent transfer activities.
- Duplicative rehabilitative therapy: When individuals receive ABA, physical, occupational, or speech therapy, the therapists should provide different treatments that reflect each therapy discipline's unique perspective on the individual's impairments and functional deficits, and not duplicate the same treatment.

## Process

### **Occupational or Physical Therapy Evaluation**

A patient with an indication for which occupational or physical therapy services may be medically necessary will be assessed by a licensed occupational or physical therapist. The occupational or physical therapy evaluation must be documented in the patient's medical record. That documentation must include the occupational or physical therapist's findings of the assessment and treatment recommendations.

### **Consideration for Enrollment in Occupational or Physical Therapy**

Enrollment in occupational or physical therapy services will be determined by the clinical judgment of a licensed occupational or physical therapist, which may include a contracted licensed occupational or physical therapist, in conjunction with the patient's PMG referring physician.

### **Plan of Care**

**Documentation Requirements:** The following care plan is required for occupational or physical therapy services:

1. Occupational or physical therapy should be provided in accordance with an ongoing, written care plan.
2. The care plan should include sufficient information to determine the medical necessity of treatment. The care plan should be specific to the diagnosis, presenting symptoms, and findings of the physical or occupational therapy evaluation. The care plan should provide for coordination of care with other members of the patient's health care team.
3. The care plan must be signed by the patient's PMG referring physician and occupational or physical therapist, when the therapist is not a KFH or PMG employee, or as required by law.
4. The care plan should include:
  - a. The diagnosis and treating diagnosis.
  - b. The date of onset or exacerbation of the disorder/diagnosis.
  - c. Specific statements of goals.
  - d. Quantitative objectives measuring current and/or age-adjusted level of functioning.
  - e. A reasonable estimate of when the goals will be reached.
  - f. The specific treatment techniques and/or exercises to be used in treatment.
  - g. The frequency and duration of treatment.
  - h. The frequency of patient re-evaluation.
5. The care plan should be ongoing and periodically updated based on reassessment by a licensed OT or PT.

6. The treatment goals and subsequent documentation of treatment results should specifically demonstrate that occupational or physical therapy services are:
  - a. contributing to reasonable functional progress made toward the goals of occupational/physical therapy, OR
  - b. medically necessary to preserve function or prevent regression.

### **Continuation of Therapy**

Continuation of therapy is based on significant, measurable improvement or is necessary to maintain skills or prevent regression in the patient's condition based on the written care plan and the clinical judgment of the treating physical or occupational therapist or healthcare professional who is competent to evaluate the specific clinical issues involved in the healthcare services requested by the provider such as those by a licensed physician or occupational or physical therapist. Regular evaluation of the patient is required to determine that continuation of therapy is medically appropriate.

### **Discontinuation of Therapy**

Occupational and physical therapy services will be discontinued when:

1. The patient has reached an age appropriate function (less than or equal to 1 SD below the mean, greater than or equal to a standard score of 85 or greater than or equal to the 16<sup>th</sup> percentile), OR
2. The patient is able to safely access their environment with supervision (i.e. walking on the grass, getting in and out of a shower, ascending and descending stairs) OR
3. the skills of a qualified provider of occupational or physical therapy services are not necessary to maintain skills or prevent regression (for example, continuation of drills, techniques and exercises by patient or caregiver after completion of medically necessary occupational or physical therapy services would be expected to preserve the patient's present level of function and prevent regression of that function) OR
4. There is an inability to benefit from therapy related to but not limited to: attention or behavior difficulties, poor attendance and/or poor compliance with home program.

Discontinuation or graduation from occupational nor physical therapy follows clinical justification and documentation of the rationale for the discontinuation/graduation from treatment.

At any time after therapy has been discontinued for the above reasons, the patient may be referred for an updated occupational or physical therapy evaluation to determine the medical necessity of resuming therapy.

These criteria address the medical necessity of existing, generally accepted services and technologies. PMG physicians can use these criteria as a resource. The criteria are not meant to replace the independent clinical judgment of the physician.

<b>SCAL Approving Bodies</b>	
Marvin L. Tan, M.D. SCAL Pediatric Behavioral Development	8/21/2017, 8/24/2018, 8/29/2019, 2/15/2021, 8/24/2021, 6/20/2022, 8/2023, 08/2024
SCAL Regional Coordinator, Autism & Developmental Disabilities	8/21/2017, 8/24/2018, 8/29/2019, 2/15/2021, 8/24/2021, 6/20/2022, 8/2023, 08/2024
John Brookey, MD William Cory, MD SCAL Health Plan UM Physician Advisors	8/21/2017, 8/24/2018, 8/29/2019, 2/15/2021, 8/24/2021, 6/20/2022, 8/2023, 08/2024
Utilization Management Steering Committee (UMSC)	8/21/2017, 8/24/2018, 8/29/2019, 2/15/2021, 08/24/202, 06/20/2022, 8/21/2017, 8/24/2018, 8/29/2019, 2/15/2021, 8/24/2021, 6/20/2022, 8/2023, 08/2024

### **UM Criteria References**

Baranek GT. Efficacy of Sensory and Motor Interventions for Children with Autism. J Autism Dev Disord. 2002;32:397-422.  
Clinical Evidence. Autism. BMJ Clin Evid 2007;12:322.

Dawson G, Waitling R. Interventions to facilitate auditory, visual, and motor integration in autism: a review of the evidence. J Autism Dev Disord. 2000;30:415-421.

ECRI Institute. Comprehensive Programs for the Treatment of Children with Autism. 1999 <http://www.ecri.org>



Escalona A, Field T, Sing-Strunck RS, Cullen C, Hartshorn K. Brief Report: Improvements in the Behavior of Children with Autism Following Massage Therapy. *J Autism Dev Disord* 2001;31:5.

Hayes Inc. Sensory Integration Therapy for Children with Learning Disabilities or Developmental Delays. 2004 Dec. <http://www.hayes.com>.

Hodgetts S, Hodgetts W. Somatosensory stimulation interventions for children with autism: literature review and clinical considerations. *Can J Occup Thera*. 2007 Dec;74(5): 393-400.

Ministry of Health New Zealand Autism Spectrum Disorder Guideline <http://www.moh.govt.nz/moh.nsf/indexmh/nz-asd-guideline-apr08>.

Scottish Intercollegiate Guidelines Network (SIGN) Assessment, diagnosis and clinical interventions for children and young people with autism spectrum disorders. A national clinical guideline. <http://www.sign.ac.uk/new.html>.

University of Western Australia Occupational Therapy Critical Appraised Topics (Sensory Integration Therapy, LEGO play, Therapy Balls). <http://www.otcats.com/topics/index.html>.

Waitling RI, Dietz J. Immediate effect of Ayres's sensory integration-based occupational therapy intervention on children with autism spectrum disorders. *Am J Occup Thr*. 2007 Sep-Oct;61(5): 574-83.

#### "Basic Activities of Daily Living" References:

- [https://www.cms.gov/Research-Statistics-Data-and-Systems/Research/MCBS/Downloads/2008\\_Appendix\\_B.pdf](https://www.cms.gov/Research-Statistics-Data-and-Systems/Research/MCBS/Downloads/2008_Appendix_B.pdf)
- <https://www.aota.org/~media/Corporate/Files/Advocacy/State/Resources/PracticeAct/Model%20Definition%20of%20OT%20Practice%20%20Adopted%2041411.pdf>
- Self-Care Measurement Tool Selection: Clearing the Confusion, OTcourses.com, By Wendy K. Anemaet, DPT, PhD, GCS, CWS, GTC, COS-C Michelle E. Moffa-Trotter, PT, GTC

#### "Episodic Care" References:

- Bailes A, Reder R, Burch C. *Peds Phys Ther* 2008; 20 :194-198, Matos M, Miller,K, Eliasson AC, Imms C. *Clin Rehab* 2007;21:47-55.
- 2012 by the Academy of Pediatric Physical Therapy, American Physical Therapy Association, 1111 N Fairfax Street, Alexandria, VA 22314-1488, [www.pediatricapta.org](http://www.pediatricapta.org).

- Bailes A, Reder R, Burch C. Development of guidelines for determining frequency of therapy services in a pediatric medical setting. *Ped Phys Ther.* 2008;20(2):194-198.
- Purtilo R. Whom to treat first and how much is enough? Ethical dilemmas that physical therapists confront as they compare individual patients' needs for treatment. *Inter J of Tech Assess in Health Care.* 1992;8:26-34.
- Centers for Disease Control and Prevention Web site. People with disabilities: living healthy. <http://www.cdc.gov/Features/Disabilities>. Accessed February 15, 2012.
- World Health Organization Web site. International classification of functioning, disability, and health. <http://www.who.int/classifications/icf/en/>. Accessed February 15, 2012.
- American Physical Therapy Association. Guide to Physical Therapist Practice. 2nd ed. Alexandria, VA: American Physical Therapy Association; 2001;81:9-744.
- American Physical Therapy Association. Standards of Practice for Physical Therapy. [http://www.apta.org/uploadedFiles/APTAorg/About\\_Us/Policies/HOD/Practice/Standards.pdf#search=%22Standards of Practice%22](http://www.apta.org/uploadedFiles/APTAorg/About_Us/Policies/HOD/Practice/Standards.pdf#search=%22Standards%20of%20Practice%22). Accessed May 4, 2012.
- Schmidt RA, Lee TD. Motor Control and Learning: A Behavioral Emphasis. 4th ed. Champaign, IL: Human Kinetics; 2005.
- Campbell S, Palisano R, Orlin M. Physical Therapy for Children. St. Louis, MO: Elsevier/Saunders; 2012:52-54.
- Zwicker J, Harris S. A reflection on motor learning theory in pediatric occupational therapy practice. *Can J Occup Ther.* 2009;76(1):29-37.
- Damiano D. Activity, activity, activity: rethinking our physical therapy approach to cerebral palsy. *Phys Ther.* 2006;86:1534-1540.
- Sullivan KJ, Katak S, Burtner PA. Motor Learning in children: feedback effects on skill acquisition. *Phys Ther.* 2008;88:720-732.
- Valvano J. Activity focused motor interventions for children with neurological conditions. *Phys Occup Ther Pediatr.* 2004;24:79-107.
- Mastos M, Miller K, Eliasson AC, Imms C. Goal directed training: linking theories of treatment to clinical practice for improved functional activities in daily life. *Clin Rehab.* 2007;21:47-55.
- Chiarello LA, O'Neil M, Dichter CG, et al. Exploring physical therapy clinical decision making for children with spastic diplegia: survey of pediatric practice. *Pediatr Phys Ther.* 2005;17(1):46-54.
- Hanft B, Feinberg E. Towards the development of a framework for determining the frequency and intensity of early intervention services. *Infant Young Child.* 1997;10(1):27-37.
- Majnemer A, Shevell MI et al. Early rehabilitation service utilization patterns in young children with developmental delays. *Child Care Health Dev.* 2005;28(1):29-37.

- Schertz M, Gordon AM. Changing the model: a call for a re-examination of intervention approaches and translational research in children with developmental disabilities. *Dev Med Child Neurol.* 2009;51(1):6-7.
- Martin L Baker R, Harvey A. A systematic review of common physiotherapy interventions in school aged children with cerebral palsy. *Phys Occup Ther Pediatr.* 2010;30(4):294-312.
- Dodd K, Taylor N, Damiano D. A systematic review of the effectiveness of strength training programs for people with cerebral palsy. *Arch Phys Med Rehab.* 2002; 83(8):1157-64.
- Dodd KJ, Shields N. A systematic review of the outcomes of cardiovascular exercise programs for people with Down syndrome. *Arch Phys Med Rehab.* 2005;86:2051-2058.
- Damiano D. Rehabilitative therapies in cerebral palsy: the good, the not as good, and the possible. *J Child Neurol.* 2009;24(9):1200-4.
- Fragala-Pinkham MA, Haley S, Goodgold S. Evaluation of a community based fitness program for children with disabilities. *Pediatr Phys Ther.* 2006;18(2):159-167.
- Bower E, McLellan DL. A randomized controlled trial of different intensities of physiotherapy and different goal setting procedures in 44 children with cerebral palsy. *Dev Med Child Neurol.* 1996;38(3):226-37.
- Bower E, Michell D, Burnett M, Campbell MJ, McLellan DL. Randomized controlled trial of physiotherapy in 56 children with cerebral palsy for 18 months. *Dev Med Child Neurol.* 2001;43(1):4-15.
- Ustad T, Sorsdahl A, Ljunggren A. Effects of intensive physiotherapy in infants newly diagnosed with cerebral palsy. *Pediatr Phys Ther.* 2009;21:140-149.
- Christiansen AS, Lange C. Intermittent versus continuous physiotherapy in children with cerebral palsy. *Dev Med Child Neurol.* 2008;50(4):290-3.
- Trahan J, Malouin F. Intermittent intensive physiotherapy in children with cerebral palsy: a pilot study. *Dev Med Child Neurol.* 2002;44(4):233-9.
- Tsorlakis N, Evaggelinou C, Grouio G, Tsorbatzoudis C. Effect of intensive neurodevelopmental treatment in gross motor function of children with cerebral palsy. *Dev Med Child Neurol.* 2004;46(11):740-745.
- Huang H, Fetters L, Hale J, McBride A. Bound for success: a systematic re-view of constraint induced movement therapy in children with CP supports improved arm and hand use. *Phys Ther.* 2009;89:1126-1141.
- Ulrich DA, Lloyd MC, Tiernan CW, Looper JE, Angulo-Barroso RM. Effects of intensity of treadmill training on developmental outcomes and stepping in infants with Down syndrome: a randomized trial. *Phys Ther.* 2008;88(1):114-122.

- Ulrich DA, Ulrich BD, Angulo-Kinzler RM, Yun J. Treadmill training of infants with Down syndrome: evidence-based developmental outcomes. *Pediatr.* 2001;108(5):E84.
- Saleh MN, Korner-Bitensky N, Snider L, et al. Actual vs. best practices for young children with cerebral palsy: a survey of paediatric occupational therapists and physical therapists in Quebec Canada. *Dev Neurorehab.* 2008;11(1):60-80.
- Montgomery P. Frequency and duration of physical therapy. *PT Mag.* 1994;42;89-91.
- Palisano R, Murr S. Intensity of therapy services: what are the considerations? *Phys Occup Ther Pediatr.* 2009;29(2):107-112.
- Ideishi R, O'Neil M, Chiarello LA, Nixon-Cave K. Perspectives of therapist's role in care coordination between medical and early intervention services. *Phys Occup Ther Pediatr.* 2010;30(1):28-42.
- [DHCS- EPSDT APL 19-010](#)